

MTK UGANDA LIMITED

GUMBORO VACCINES (CEVAC® GUMBO L & CEVAC® IBD L)

Vaccine	Composition	Indications	Precautions & Contraindications	Route	Dosage and administration
CEVAC® GUMBO L	LIBD Strain of infectious Bursal disease virus live freeze dried form	Vaccination of healthy birds against IBD	<ul style="list-style-type: none"> • Storage is at 2°C - 8° C • Protected from light • All birds in the flock must be vaccinated at the same time • Administer at coolest time of the day (preferably morning hours) • Vaccinate healthy birds. • Should not be mixed with other vaccines • Dispose off all opened vaccine containers remaining after vaccination. • Don't use DISINFECTANT in drinking water 48hrs prior to & 24 hrs after vaccination. • Withhold water from birds on average of 1½-2 hrs before administration. 	Oral (Drinking water)	<p>Broilers; Btn 14-21 days Pullets; Btn 20-28 days</p> <p>Chicks from Non vaccinated parent stock; Broilers; 7days Pullets;14 days</p> <p>NB; Pullets should be vaccinated twice following a 6 day interval.</p>
CEVAC® IBD L	Live freeze dried vaccine, Winterfield 2512 G-61 strain, for the active immunization of chickens against IBD	Vaccination of healthy birds against IBD	<ul style="list-style-type: none"> • Storage is at 2°C - 8° C • Protected from light • All birds in the flock must be vaccinated at the same time • Administer at coolest time of the day (preferably morning hours) • Vaccinate healthy birds. • Should not be mixed with other vaccines • Dispose off all opened vaccine containers remaining after 	Oral (Drinking water)	<p>Broilers; 10-18 days Pullets; 16-26 days allowing six day interval.</p>

			<p>vaccination.</p> <ul style="list-style-type: none"> • Don't use DISINFECTANT in drinking water 48hrs prior to & 24 hrs after vaccination. • Withhold water from birds on average of 1½-2 hrs before administration. • Do not use water containing high concentrations of metal ions. • Preserving vaccine activity, neutralize the chlorine water with 2.5g of skimmed milk powder per liter of water. 		
--	--	--	---	--	--

VACCINE MIXING RATES

1. CEVAC GUMBO L

Age	7-12 days	13-18 days	19-28 days
Minimum amount of water for 1000 birds	10 L	15-20 L	25-30 L

2. CEVAC IBD L

Age	10-12 days	13-18 days	19-26 days
Minimum amount of water for 1000 birds	10 L	15 L	25-30 L